

Conditional parameter estimates from Mixed Logit models: distributional assumptions and a free software tool

Stephane Hess*

July 1, 2010

Abstract

A number of authors have discussed the possible advantages of conditioning parameter distributions on observed choices when working with Mixed Multinomial Logit models. However, the number of applications is still relatively small, partly due to a limited implementation in available software. To address this situation, the present paper discusses the development of a freeware software tool that allows users to compute conditional distributions independently of the software used during model estimation. Additionally, the paper looks at what impact assumptions made for the unconditional distributions have on the results obtained with conditional distributions. Here, an application using stated choice data collected in Denmark shows that while the move from unconditional to conditional distributions possibly brings results closer together, some discrepancies do remain.

KEYWORDS: mixed logit; discrete choice; conditional distributions; taste heterogeneity

1 Introduction

The random coefficients formulation of the Mixed Multinomial Logit (MMNL) model (cf. [Revelt and Train, 1998](#); [Train, 1998](#); [McFadden and Train, 2000](#); [Hensher and Greene, 2003](#); [Train, 2003](#)) is fast becoming one of the most widely used econometric structures for the analysis of choice behaviour. The main advantage of the MMNL model over its more simplistic closed-form counterparts is that it

*Institute for Transport Studies, University of Leeds, s.hess@its.leeds.ac.uk, Tel: +44 (0)113 34 36611, Fax: +44 (0)113 343 5334

allows for a relaxation of the assumption of constant marginal utility coefficients across individuals.

The MMNL model accommodates taste heterogeneity by allowing marginal utility coefficients to be distributed randomly across respondents. A major issue in this context is the choice of an appropriate mixing distribution in the absence of information on the actual shape of that distribution in the sample population (see for example [Hensher and Greene 2003](#), [Hess et al. 2005](#) and [Fosgerau 2006](#)). The vast majority of MMNL applications make use of the Normal distribution. Here, problems can arise due to the unbounded nature of the distribution, as well as due to its symmetry assumption. These can lead to issues with sign violations and biased mean values respectively. Indeed, it is in such situations not clear whether the findings actually reflect real sensitivities present in the data or are simply a result of the distributional assumptions. A possible solution is to use more flexible distributions, not making a strict symmetry assumption, while also allowing for the estimation (rather than imposition) of bounds to either side. Examples of such distributions include the Johnson S_B , discussed in detail in a MMNL context by [Train and Sonnier \(2005\)](#).

A serious problem is that models making use of these more advanced distributions are considerably more difficult to estimate than their counterparts relying on more restrictive distributions, often leading to issues with convergence or parameter significance. Another major issue with the MMNL model is that while it allows the user to accommodate random taste heterogeneity in the sample population, it does not directly provide any information on the likely location of a given respondent on this distribution. However, simply knowing that a coefficient varies across respondents is only of limited practical use.

An obvious way of dealing with this second issue is to move from the unconditional (i.e. sample population level) distribution to a conditional distribution. This equates to inferring the likely position of each sampled individual on the distribution of sensitivities (cf. [Revelt and Train, 1999](#); [Train, 2003](#); [Sillano and Ortúzar, 2004](#); [Greene et al., 2005](#)).

Let β give a vector of taste coefficients that are jointly distributed according to $f(\beta | \Omega)$, where Ω is a vector of distributional parameters that is to be estimated from the data. Let Y_n give the sequence of observed choices for respondent n (which could be a single choice), and let $L(Y_n | \beta)$ give the probability of observing this sequence of choices with a specific value for the vector β . Then it can be seen that the probability of observing the specific value of β given the choices of respondent n is equal to:

$$L(\beta | Y_n) = \frac{L(Y_n | \beta) f(\beta | \Omega)}{\int_{\beta} L(Y_n | \beta) f(\beta | \Omega) d\beta} \quad (1)$$

The integral in the denominator of Equation 1 does not have a closed form solution, such that its value needs to be approximated by simulation. This is a simple (albeit numerically expensive) process, with as an example the mean for the conditional distribution for respondent n being given by:

$$\widehat{\beta}_n = \frac{\sum_{r=1}^R [L(Y_n | \beta_r) \beta_r]}{\sum_{r=1}^R L(Y_n | \beta_r)}, \quad (2)$$

where β_r with $r = 1, \dots, R$ are independent multi-dimensional draws¹ with equal weight from $f(\beta | \Omega)$ at the estimated values for Ω . Here, $\widehat{\beta}_n$ gives the most likely value for the various marginal utility coefficients, conditional on the choices observed for respondent n .

It is important to stress that the conditional estimates for each respondent follow themselves a random distribution, and that the output from Equation 2 simply gives the expected value of this distribution. As such, a distribution of the output from Equation 2 across respondents should not be seen as a conditional distribution of a taste coefficient across respondents, but rather a distribution of the means of the conditional distributions (or conditional means) across respondents. Here, it is similarly possible to produce a measure of the conditional standard deviation, given by:

$$\widetilde{\beta}_n = \sqrt{\frac{\sum_{r=1}^R [L(Y_n | \beta_r) (\beta_r - \widehat{\beta}_n)^2]}{\sum_{r=1}^R L(Y_n | \beta_r)}}, \quad (3)$$

with $\widehat{\beta}_n$ taken from Equation 2.

Obtaining information on the likely location of a given respondent on the distribution of tastes across the sample population can be a great asset for various reasons. Here, [Greene et al. \(2005\)](#) and [Hess \(2007\)](#) amongst others show that when using conditional means, issues with counter-intuitively signed coefficients are largely avoided. However, these applications fail to recognise that the conditional values themselves follow a distribution, and the ratio of the conditional mean time and cost sensitivities for an individual is as a consequence not the same as the mean of the ratio of the individual specific conditional distributions for the time and cost sensitivities. Other applications have been more concerned with making use of the conditional estimates for individual coefficients with a view to informing various classification approaches. Here, one application

¹The term *independent* relates to independence across different multivariate draws, where the individual multivariate draws allow for correlation between univariate draws.

comes in attempts to retrieve individual specific information processing strategies (cf. [Hess and Hensher, 2008](#)), while [Campbell and Hess \(2009\)](#) have recently explored the use of conditional parameter distributions in the process of identifying respondents with extreme sensitivities, i.e. outliers, in data used for discrete choice models. Finally, as discussed for example by [Train \(2003\)](#), information on individual specific distributions can be used in posterior analyses (e.g. cluster analysis) that identify different segments of respondents and link the heterogeneity to socio-demographic attributes. This can once again inform re-specification of the model, this time with different or greater segmentation. However, as noted by [Train \(2003\)](#), this approach is only applicable if the conditional means themselves account for a sufficiently large share of the heterogeneity in the sample distribution.

What has received relatively little attention in the existing literature is the potential impacts of the unconditional distributional assumptions on the shape of the conditional distributions. As discussed by [Train \(2003\)](#), the combination of respondent-specific distributions across the sample yields the sample distribution. However, the interest is in the individual-specific distributions, and in particular in many cases the conditional means, giving the most likely position of each individual in the sample distribution. Here, if the conditional distributions could be shown to be relatively independent on the assumptions made for the former, analysts could rely on easier to use unconditional distributions (e.g. Normal) if the aim is to make use of the means of these conditional distributions. If any out of sample prediction work was planned, then conditional distributions are clearly not applicable. However, if conditional distributions are indeed less affected by distributional assumptions, and give a better indication of the actual *true* distribution, then they can potentially be of use in informing a better choice of distributional assumptions for a *revised* model.

Another issue limiting the use of conditional parameter distributions is the lack of available software, with only NLogit ([Econometric Software, 2007](#)) giving users the possibility of producing conditional parameters. Despite the popularity of NLogit, many analysts rely on other packages for MMNL analyses, notably Biogeme ([Bierlaire, 2005](#)) or purpose written code.

Given the above discussion, the aims of this paper are twofold. Firstly, the paper presents a freeware software tool that is able to produce conditional parameter estimates for a range of different model specifications (i.e. distributional assumptions), independently of the software package used in model estimation. Secondly, we present an application that discusses the impacts of assumptions on the shape of the unconditional distribution on the resulting shape of the conditional distribution.

The remainder of this paper is organised as follows. The free software tool is

described in Section 2. Section 3 presents the empirical application comparing different distributions. Finally, Section 4 presents the conclusions of the paper.

2 Free software tool

The free software was produced using Matlab and is available for download from the author’s website². The programme consists of a standalone executable³ along with a spreadsheet tool used to generate the input files for the Matlab programme. We will now look at these two components in turn.

The macro-driven spreadsheet initialises to the situation shown in the first half of Figure 1. Here, the user is required to specify the number of coefficients, observations, alternatives and respondents. Additionally, an output directory needs to be specified, alongside a name for the model. In the second half of Figure 1, we have used the settings from the empirical application in Section 3, and have specified a *conditionals* subdirectory of the C drive along with naming our model *example*.




Figure 1: Free software tool: initial settings (before and after)

The next step is for the user to press the *Generate fields* button. This produces fields in which the user needs to enter respondent identification numbers, choice

² www.stephanehess.me.uk

³For users without a Matlab installation, the programme requires the installation of libraries that are similarly available for download from the author’s website.




Figure 2: Free software tool: data entry stage (before and after)

indicators and the attributes of the various alternatives⁴. Additionally, the software generates a number of fields for each of the coefficients where the user needs to make a choice of distribution and enter the estimated parameters. A pop-up window explains the various settings to the user, and some explanations are also provided later in the paper in the context of the empirical application.

The situation after pressing the *Generate fields* button is illustrated in the first half of Figure 2. The second half shows the situation after the settings have been entered. Here, the first parameter is a constant, fixed at 0.37, while the travel time and travel cost coefficients follow Normal distributions, with means and standard deviations given by the α and γ parameters, taken from the application in Section 3. The spreadsheet is limited to coefficients using independent Normal, Uniform, symmetrical Triangular, Lognormal and Johnson S_B distribu-

⁴Here, the software is limited to a linear in parameters specification, but the user can work around this to include interactions with socio-demographic variables by making some attributes specific to given socio-economic subgroups.

tions. These limitations however do not apply to the Matlab programme, and the user can also directly generate input files making use of other distributions, including multi-variate ones (e.g. multivariate Normal with Cholesky transformation).

The figure shows all 8 choices for the first respondent, along with the first 3 choices for the second respondent. The attribute levels are entered in block for each alternative, using the same ordering as in the section specifying the coefficients. In the present example, the first attribute is a dummy variable used for the first alternative, with the second and third giving travel cost and travel time respectively⁵.

The next two steps require the user to first run a check on the entered values (the *Check input* button) before using the *Produce outputs* button to generate the input files for the Matlab programme. This latter button leads to the generation of three separate files, one containing the data, one containing the draws to be used for the various coefficients and one containing the settings of the problem in terms of coefficients, observations, alternatives and respondents⁶. The file names are based on the name specified by the user in cell G3.

After completing the generation of the input files, the user is now ready to proceed to the Matlab tool for the computation of the conditional parameter estimates. After launching the executable, the user is prompted to enter the name of the model, as specified in cell G3 in the spreadsheet tool. The remainder of the process requires no user input and is illustrated in Figure 3. As a first step, the software provides an overview of the data in terms of choices, respondents and coefficients. It then gives some summary statistics for the simulated draws for the various coefficients. Finally, it shows the unconditional log-likelihood⁷, the log-likelihood calculated with the conditional distributions for each respondent (i.e. assigning the conditional weights from Equation 1 to the individual draws), and the log-likelihood calculated with the means of the conditional distributions for each respondent (i.e. making use of just one value for each respondent).

The software produces two output files⁸. The actual conditionals are saved in a file that contains two columns for each coefficient, namely the mean and standard deviation of the conditional distribution for each individual. Additionally,

⁵Travel cost is given in øre, where one Danish Krone is equal to 100 øre, with travel time given in minutes.

⁶If the inputs are generated manually, i.e. not making use of the spreadsheet, the user needs to similarly generate the appropriate three files.

⁷Here, some discrepancies are possible when compared to the log-likelihood produced during estimation due to the use of different random draws. As an example, the unconditional and conditional log-likelihood values in Figure 3 differ slightly from those in Table 1 and Table 2.

⁸The model name used in the input is used as the basis of the filenames for the output.

```

C:\conditionals\conditionals.exe
#####
# Calculation of conditional parameters estimates from Mixed Logit models #
#   Stéphane Hess, Institute for Transport Studies, University of Leeds   #
#   s.hess@its.leeds.ac.uk; www.stephanehess.me.uk                       #
#####
Please enter the name of your model, as specified in spreadsheet cell G3: example
Reading in data
Completed reading inputs
Number of included choices : 1767
Number of individuals : 230
Number of coefficients : 3

Summary for simulated coefficients
Coefficient 1: Mean=0.37, std.dev.=7.777e-014
Coefficient 2: Mean=-0.003559, std.dev.=0.001718
Coefficient 3: Mean=-0.1451, std.dev.=0.1795

Starting calculations
Calculations completed
Log-likelihood with parameter estimates: -1039.304
Log-likelihood with full conditional distribution: -863.0299
Log-likelihood at conditional means: -822.0189

Saving outputs

Main outputs saved in file example_conditionals.dat
Format of file:
Column 1: respondent id
Column 2: mean for conditional distribution for first coefficient
Column 3: std. dev. for conditional distribution for first coefficient
etc.

Respondent-specific weights for conditional draws saved in file example_weights.dat
Press ENTER to exit

```

Figure 3: Free software tool: calculation of conditionals

the software produces a file that contains for each individual the weights for the 10,000 draws used as the input. The weights for a given draw from β , say β_p , is given by $\frac{L(Y_n|\beta_p)}{\sum_{r=1}^R L(Y_n|\beta_r)}$. On the basis of these weights, it is then possible to produce draws from the conditional distribution for each respondent.

3 Empirical application

This section presents the findings of a brief application discussing the impacts of unconditional distributions on the shape of the conditional distributions. Model estimation was carried out in Biogeme (Bierlaire, 2005), making use of 500 Halton draws per individual, and the above discussed tool was used for the generation of the conditional parameter estimates. To ensure consistency of the results, the unconditional Biogeme estimates and conditional parameter estimates from the software tool discussed in this paper were compared to results obtained in NLogit (Econometric Software, 2007), with the exception of the Johnson S_B distribution,

which is only implemented with some parameter restrictions in NLogit.

3.1 Data

The analysis makes use of stated choice (SC) data collected for the DATIV study carried out in Denmark in 2004 (cf. [Burge and Rohr, 2004](#)). For this survey, a binary unlabelled route choice experiment was used, with two attributes, travel time and travel cost describing the alternatives. For the present analysis, we make use of 1,767 observations collected from 230 respondents.

3.2 Model specification

Across all models used, a constant was associated with the first alternative (δ_1), and the travel time (β_{TT}) and travel cost (β_{TC}) coefficients were interacted linearly with the associated attributes.

Depending on the specification of taste heterogeneity, up to four parameters (a , b , α and γ) were estimated for each coefficient, where, in the context of this illustrative example, univariate distributions were used. We will now look at the various models in turn.

MNL: In the MNL model, a point estimate (α) was estimated for both coefficients.

Uniform: In the MMNL model making use of a Uniform distribution, a left boundary (a) was estimated along with a positive range parameter (b)⁹.

Triangular: In the MMNL model making use of a Triangular distribution, a left boundary (a) was estimated along with a positive range parameter (b)¹⁰. The distribution was constrained to be symmetrical, with the mean, median and mode being equal to $\frac{a+b}{2}$.

Normal: In the MMNL model making use of a Normal distribution, the mean is given by α , with the standard deviation given by γ

Lognormal: In the MMNL model making use of a Lognormal distribution, α and γ give the mean and standard deviation respectively for the underlying Normal distribution. The offset parameter a is either positive or negative, and b is a direction parameter, which is equal to -1 for both coefficients, resulting in a tail towards minus infinity.

⁹Note that these parameters were obtained as transformations of the original Biogeme estimates which are for the mean and half the spread.

¹⁰Once again, transformations of Biogeme estimates were used.

Johnson S_B : In the MMNL model making use of a Johnson S_B distribution, α and γ again give the mean and standard deviation respectively for the underlying Normal distribution. The offset parameter a is again either positive or negative, and b is a positive range parameter.

3.3 Estimation results

The estimation results are summarised in Table 1. Here, we can see that all five MMNL specifications lead to significant gains in model fit over the MNL model, highlighting the presence of significant levels of taste heterogeneity in the sample. In terms of the adjusted ρ^2 measure, the best performance is obtained by the Johnson S_B distribution ahead of the Lognormal distribution, with the symmetrical Triangular giving the poorest fit to the data. The three symmetrical distributions produce significant probabilities of positive coefficient values, especially for β_{TT} , where these results can be directly linked to the distributional assumptions (cf. Hess et al., 2005). Both the Lognormal and the Johnson S_B distributions indicate that the domain of the distribution for the two taste coefficients should be entirely negative. Indeed, for the Lognormal distribution, both offset parameters are negative (albeit no different from zero), while for the Johnson S_B distribution, the offset and range parameters are such that the upper limit for both coefficients is below zero. Before moving on, it should also be noted that for the Johnson S_B distribution, the second shape parameter (γ) is only significantly different from zero at the 89% level for β_{TC} , while for β_{TT} , neither shape parameter is significant at any reasonable level of confidence. This is an illustration of the difficulties of estimating parameters for this complex distribution.

3.4 Conditional model results

After estimation of the five MMNL models, the tool developed in Section 2 was used to produce means and standard deviations for the conditional distributions for each respondent. As a first illustration of the additional information gained from this, Table 2 compares $LL(\Omega)$, the log-likelihood measures obtained during estimation (i.e. using the sample distribution parameters), to $LL(\hat{\Omega}_n)$, the log-likelihood measures obtained using the individual-specific conditional distributions, and $LL(\hat{\beta}_n)$, the log-likelihood measures obtained using the means of the conditional distributions.

For each of the five distributions, we observe dramatically better fit when making use of the conditional distributions as opposed to making use of the unconditional distributions. This is to be expected as we now calculate the choice

Table 1: Estimation results

	MNL	Uniform	Triangular	Normal	Lognormal	Johnson S_B
Final LL	-1,096.64	-1,038.06	-1,042.58	-1,038.43	-1,033.12	-1,027.07
Parameters	3	5	5	5	7	9
adj. ρ^2	0.1020	0.1480	0.1447	0.1480	0.1510	0.1540

	est.	t-rat.	est.	t-rat.	est.	t-rat.	est.	t-rat.	est.	t-rat.	est.	t-rat.
δ_1	0.3020	5.92	0.3680	6.33	0.3710	6.39	0.3700	6.35	0.3630	6.30	0.366	6.32
$\beta_{TC} (a)$	-	-	-0.0074	-7.50	-0.0100	-8.99	-	-	-0.0001	-0.16	-0.0148	-3.62
$\beta_{TC} (b)$	-	-	0.0070	5.94	0.0123	-7.94	-	-	-1	-	0.013	3.14
$\beta_{TC} (\alpha)$	-0.0017	-10.44	-	-	-	-	-0.0036	-9.49	-5.7800	-13.97	3.41	2.72
$\beta_{TC} (\gamma)$	-	-	-	-	-	-	0.0017	5.64	0.7370	2.86	4.8	1.6
$\beta_{TC} (\% > 0)$	-	-	0.00%		6.45%		1.89%		0.00%		0.00%	
$\beta_{TT} (a)$	-0.4370	-6.59	-0.5816	-7.49	-0.5816	-7.49	-	-	-0.0250	-0.41	-0.307	-7.21
$\beta_{TT} (b)$	0.5440	4.51	0.7640	-6.20	0.7640	-6.20	-	-	-1	-	0.249	5.39
$\beta_{TT} (\alpha)$	-0.0829	-6.37	-	-	-	-	-0.1470	-5.91	-2.2500	-3.91	-3.75	-0.59
$\beta_{TT} (\gamma)$	-	-	-	-	-	-	0.1780	6.60	0.8890	3.26	13.9	0.73
$\beta_{TT} (\% > 0)$	-	-	19.67%		10.86%		20.44%		0.00%		0.00%	

Table 2: Log-likelihood at convergence and using conditional distributions, and means of conditional distributions

	$LL(\Omega)$	$LL(\Omega_n)$	$LL(\hat{\beta}_n)$
Uniform	-1,038.06	-862.52	-820.46
Symmetrical triangular	-1,042.58	-853.24	-807.86
Normal	-1,038.43	-863.03	-822.02
Lognormal	-1,033.12	-859.11	-821.99
Johnson S_B	-1,027.07	-863.74	-808.76

probabilities for each individual by drawing from a distribution for the random coefficients that is more likely to be the *true* distribution for that respondent. Further increases are obtained when relying solely on the means of the conditional distributions, i.e. using for each respondent the most likely values for the two coefficients.

Surprisingly, the best performance for the two conditional log-likelihood measures is obtained by the Triangular distribution, even though it produced the lowest log-likelihood measure in estimation (cf. Table 1). Furthermore, while the Johnson S_B distribution obtained the best fit in estimation, it produces the lowest measure for $LL(\hat{\Omega}_n)$, although, alongside with the Triangular distribution, it then produces the best performance when working with the means of the conditional distributions, i.e. $LL(\hat{\beta}_n)$. This could suggest some differences in how well various distributions can be used to infer individual specific distributions post estimation. However, it is not entirely clear what could be causing this interesting finding. A possible explanation could have been discrepancies between the unconditional distributions and the aggregated (over respondents) conditional distributions. As mentioned earlier, the aggregated conditional distributions should be equal to the unconditional distributions. If for example, the Uniform, Normal, Lognormal, and Johnson S_B distributions had failed that test, this would have indicated that they are incorrect distributions for the present application, unlike the Triangular¹¹. However, all five distributions passed the test, thus not indicating any inherent problems with one of the distributions, but rather supporting the above point about differences across distributions in the impact of the distributional assumptions on the conditional distributions.

As a first step in our comparison of the results across models, we look solely at the conditional means for the two coefficients, i.e. the most likely values for the

¹¹The author is grateful to Kenneth Train for this suggestion.

time and cost coefficients for each respondent. Empirical distribution functions for the conditional means for the two coefficients are shown in Figure 4, alongside the unconditional distributions. Here, we can see that for all five models, the distribution of the conditional means has a narrower range than the unconditional distribution, where sign violations have also almost completely disappeared, repeating earlier results by [Greene et al. \(2005\)](#). For the cost coefficient, the distribution has a significantly longer tail for the Lognormal and Johnson S_B distributions, while, for the time coefficient, the ranges are more comparable.

So far, we have looked at the empirical distribution function for the conditional means across respondents, and have compared these distributions across the five models. What is also interesting is to look at the results for each respondent separately and compare these across respondents. This is the approach taken in Figure 5, where, for clarity, the respondents are sorted according to the conditional means produced by the Uniform model. This approach was used solely with a view to analysing the stability across distributions in the *ordering* of conditional means, and as such, the specific choice of a base distribution should have only limited impact. Here, we can see that for β_{TC} , the results are relatively stable for respondents with low cost sensitivity, while, for respondents with high cost sensitivity, the sensitivities produced by the Lognormal and especially the Johnson S_B distribution are more extreme. For β_{TT} , the results are more stable for respondents with high time sensitivity with the exception of the Lognormal distribution, while there is now a greater discrepancy across distributions when looking at respondents with low time sensitivity.

As a next step, we conduct an analysis similar to that reported in [Train \(2003, section 11.6.2.\)](#), by comparing measures of mean and standard deviation for the sample level distribution and for the distribution of the conditional means across respondents. The interest in this process is to see what share of the sample distribution variance is explained through making use of the conditional means only, hence disregarding any variation around these conditional means. The results of this process are summarised in Table 3, where μ_β and σ_β refer to the means and standard deviations of the estimated sample level distributions for the two coefficients, and μ_{μ_n} and σ_{μ_n} refer to the means and standard deviations for the distribution of the conditional means across respondents.

As expected for a well specified and consistently estimated model, there are only very small differences between the sample distribution means (μ_β) and the means of the distribution of conditional means (μ_{μ_n}). Turning our attention to the estimated standard deviations and the standard deviations for the distribution of conditional means across respondents, we observe major differences, with σ_{μ_n} always being below σ_β . This implies that not accounting for the heterogeneity in the conditional distributions gives only a partial representation of the




Figure 4: Unconditional distributions and distribution of conditional means for β_{TC} and β_{TT}




Figure 5: Distribution of conditional means for β_{TC} and β_{TT} , sorted by values obtained with Uniform distribution

Table 3: Mean and standard deviation for sample level distributions and for distribution of conditional means

	β_{TC}			β_{TT}		
	μ_β	μ_{μ_n}	$\frac{\mu_{\mu_n}}{\mu_\beta}$	μ_β	μ_{μ_n}	$\frac{\mu_{\mu_n}}{\mu_\beta}$
Uniform	-0.0039	-0.0038	0.98	-0.1591	-0.1671	1.05
Symmetrical triangular	-0.0039	-0.0041	1.05	-0.1971	-0.1902	0.96
Normal	-0.0036	-0.0035	0.99	-0.1470	-0.1528	1.04
Logormal	-0.0042	-0.0042	1.01	-0.1815	-0.1797	0.99
Johnson S_B	-0.0051	-0.0052	1.02	-0.2088	-0.2026	0.97

	β_{TC}			β_{TT}		
	σ_β	σ_{μ_n}	$\frac{\sigma_{\mu_n}}{\sigma_\beta}$	σ_β	σ_{μ_n}	$\frac{\sigma_{\mu_n}}{\sigma_\beta}$
Uniform	0.0020	0.0012	0.61	0.1563	0.0952	0.61
Symmetrical triangular	0.0025	0.0015	0.61	0.1565	0.0891	0.57
Normal	0.0017	0.0009	0.54	0.1780	0.1140	0.64
Logormal	0.0034	0.0020	0.58	0.1717	0.0890	0.52
Johnson S_B	0.0046	0.0035	0.75	0.1145	0.0549	0.48

heterogeneity in the data. There are some differences across distributions, but on average, we can see that making use of only the conditional means, we recover just over half the sample level heterogeneity. This means that the share of the differences across respondents that is captured through making use of the means of the conditional distribution is in this case potentially large enough to allow us to use this information to distinguish between respondents, for example in cluster analysis. However, it also means that the individual specific coefficient values are not known with certainty (in which case the standard deviation of the distribution of conditional means would be equal to the sample level standard deviation), and that it would hence not be adequate to make use of these conditional means for example in the calculation of willingness to pay measures.

After first concentrating entirely on the conditional means, we now incorporate the uncertainty in the individual specific conditional distributions. As highlighted previously, there is, for each individual, heterogeneity around the conditional mean, and the aggregation of the conditional distributions across individuals yields the sample level distribution. However, the degree of heterogeneity for individual conditional distributions should on average be expected to be lower than the sample level degree of heterogeneity, as this latter measure also incorporates heterogeneity in the mean values across respondents. We un-

Table 4: Degree of heterogeneity expressed as coefficient of variation

Distribution	β_{TC}			β_{TT}		
	cv	$\widehat{\mu}(cv)$	$\widehat{\mu}_{0.5}(cv)$	cv	$\widehat{\mu}(cv)$	$\widehat{\mu}_{0.5}(cv)$
Uniform	0.52	0.45	0.44	0.98	1.33	0.80
Symmetrical triangular	0.64	0.73	0.45	0.79	0.88	0.66
Normal	0.48	0.45	0.39	1.21	184.11	0.83
Lognormal	0.82	0.55	0.55	0.95	0.62	0.62
Johnson S_B	0.91	0.49	0.48	0.55	0.54	0.56

dertake this comparison by calculating the coefficient of variation for β_{TC} and β_{TT} for each respondent. The results of this process are summarised in Table 4. Here, we show three measures for each coefficient, namely the coefficient of variation of the unconditional distribution (cv), the mean across respondents of the coefficient of variation for the conditional distributions ($\widehat{\mu}(cv)$) and the corresponding median ($\widehat{\mu}_{0.5}(cv)$). The reason for including the latter is that it is less sensitive to outliers. Looking first at the travel cost coefficient, we see significant differences across estimated distributions, with much higher variation for the Lognormal and Johnson S_B distributions. When looking at the conditional distributions, especially in terms of the median, the results are far more stable across the five distributions. Turning our attention to the travel time coefficient, the main outliers when looking at the unconditional distributions are the Normal (high) and the Johnson S_B (low). With the conditional distributions, the results are again more similar, though only when working with the median given the huge outliers with the Normal distribution that have a major impact on the mean. Comparing the median of the coefficient of variation for the conditional distribution ($\widehat{\mu}_{0.5}(cv)$) to the coefficient of variation for the sample level distribution (cv), we observe, with the exception of the Johnson S_B for β_{TT} , a reduction in the degree of heterogeneity. This is consistent with the observation in Table 3 that part of the heterogeneity is captured in the variation across respondents in the conditional means, where this share of the heterogeneity is lowest for the Johnson S_B for β_{TT} .

As a final step, Figure 6 shows a graphical analysis of the results summarised in Table 4 where we again sort the result by respondent according to the measures obtained with the Uniform models. Here, we can see significant differences across the five distributions especially for respondents with high degrees of uncertainty in the conditional distributions, with the biggest outliers arising from the Triangular and Normal models. This also shows the large differences across respondents which could not be reflected in the results in Table 4. The same applies for





Figure 6: Coefficient of variation for conditional distributions, sorted by respondent

the time coefficient, with even bigger discrepancies for respondents with high degrees of uncertainty. In contrast with Figure 5, this shows that while the results are quite stable across distributions in terms of the means of the conditional distributions, i.e. the most likely location of each respondent on the sample level distribution, differences arise in the variation around these mean levels. This is also consistent with an observation that can be made for the unconditional distributions on the basis of Table 3, namely that while the mean values are relatively stable across the five distributions, there are much larger differences when it comes to the retrieved degree of heterogeneity.

4 Summary and conclusions

This paper has discussed the issue of the computation of conditional distributions for coefficients estimated using continuous Mixed Multinomial Logit models. While this topic has been looked at at length by various authors, as discussed in Section 1, the number of applications making use of conditional distributions is still relatively limited. This paper has identified the lack of available software (other than NLogit, [Econometric Software 2007](#)) as one reason for this and has consequently discussed the development of a freeware software tool that allows users to compute conditional distributions from any choice of unconditional distributions¹², independently of the software used during model estimation.

The paper has also looked at an additional issue in this area, namely the impact of assumptions made for the unconditional distributions on the shape of the conditional distributions. Here, an application using stated choice data collected in Denmark has shown that while the move from unconditional to conditional distributions potentially brings results closer together (notably in terms of the conditional means), some discrepancies do remain. In this context, further work is required, notably a large scale study making use of simulated data with various underlying *true* distributions. It is also important to acknowledge a limitation of the present study in that it does not take into account the sampling distribution of the parameters of the underlying distribution, a further development in the context of conditional distributions, discussed by [Train \(2003, section 11.3\)](#).

Acknowledgements

This paper is based on work funded by a University of Leeds Knowledge Transfer grant, with further developments undertaken thanks to a Leverhulme Early

¹²Noting again that the limitations in terms of distributional assumptions only apply to the spreadsheet tool, not the Matlab programme.

Career Fellowship. The inspiration for the empirical analysis stems from earlier work conducted with John Rose in the Institute of Transport and Logistics Studies at the University of Sydney. The author would also like to thank Richard Connors for coding suggestions, and Kenneth Train for useful feedback.

References

- Bierlaire, M., 2005. An introduction to BIOGEME Version 1.4. biogeme.epfl.ch.
- Burge, P., Rohr, C., 2004. DATIV: SP Design: Proposed approach for pilot survey. Tetra-Plan in cooperation with RAND Europe and Gallup A/S.
- Campbell, D., Hess, S., 2009. Outlying sensitivities in discrete choice data: causes, consequences and remedies. paper presented at the European Transport Conference, Noordwijkerhout, The Netherlands.
- Econometric Software, 2007. Nlogit 4.0. Econometric Software, New York and Sydney.
- Fosgerau, M., 2006. Investigating the distribution of the value of travel time savings. *Transportation Research Part B* 40 (8), 688–707.
- Greene, W. H., Hensher, D. A., Rose, J. M., 2005. Using classical simulation-based estimators to estimate individual wtp values. In: Scarpa, R., Alberini, A. (Eds.), *Applications of Simulation Methods in Environmental and Resource Economics*. Springer Publisher, Dordrecht, Ch. 2, pp. 17–33.
- Hensher, D. A., Greene, W. H., 2003. The Mixed Logit Model: The State of Practice. *Transportation* 30 (2), 133–176.
- Hess, S., 2007. Posterior analysis of random taste coefficients in air travel choice behaviour modelling. *Journal of Air Transport Management* 13 (4), 203–212.
- Hess, S., Bierlaire, M., Polak, J. W., 2005. Estimation of value of travel-time savings using mixed logit models. *Transportation Research Part A* 39 (2-3), 221–236.
- Hess, S., Hensher, D. A., 2008. Using conditioning on observed choices to retrieve individual-specific attribute processing strategies. paper submitted to *Transportation Research B*.
- McFadden, D., Train, K., 2000. Mixed MNL Models for discrete response. *Journal of Applied Econometrics* 15, 447–470.

- Revelt, D., Train, K., 1998. Mixed Logit with repeated choices: households' choices of appliance efficiency level. *Review of Economics and Statistics* 80 (4), 647–657.
- Revelt, D., Train, K., 1999. Customer-specific taste parameters and Mixed Logit: Households' choice of electricity supplier. Working Paper No. E00-274. Department of Economics, University of California, Berkeley, CA.
- Sillano, M., Ortúzar, J. de D., 2004. Willingness-to-pay estimation from mixed logit models: some new evidence. *Environment & Planning A* 37 (3), 525–550.
- Train, K., 1998. Recreation demand models with taste differences over people. *Land Economics* 74, 185–194.
- Train, K., 2003. *Discrete Choice Methods with Simulation*. Cambridge University Press, Cambridge, MA.
- Train, K., Sonnier, G., 2005. Mixed logit with bounded distributions of correlated partworths. In: Scarpa, R., Alberini, A. (Eds.), *Applications of Simulation Methods in Environmental and Resource Economics*. Springer Publisher, Dordrecht, The Netherlands, Ch. 7, pp. 117–134.